

AD 673 021

ELECTROPHORESIS OF SERUM PROTEINS OF SPOTTED FEVER

Yu. A. Krichevskii

Army Biological Laboratories
Frederick, Maryland

1 July 1963

AD 673021


TRANSLATION NO. 753

DATE: 1 July 1963

DDC AVAILABILITY NOTICE

Qualified requestors may obtain copies of this document from DDC.

This publication has been translated from the open literature and is available to the general public. Non-DOD agencies may purchase this publication from the Clearinghouse for Federal Scientific and Technical Information, U. S. Department of Commerce, Springfield, Va.



Reproduced by the
CLEARINGHOUSE
for Federal Scientific & Technical
Information Springfield Va 22151

DEPARTMENT OF THE ARMY
Fort Detrick
Frederick, Maryland

This document has been approved
for public release and sale; its
distribution is unlimited.

#753
g.1
Laboratornoye Delo (Russian Journal #7, 1961, pp 19-20).

Translated by Sp.6 Charles T. Ostertag Jr.

Electrophoresis of Serum Proteins of Spotted Fever. Yu. A. Krichevskiy.
Infectious Disease Clinic (Chief lecturer G. A. Fridman) of the Kharkov Medical
Institute.

In 1957 we undertook a dynamic study of the serum proteins of spotted fever
by a method of electrophoresis on paper with the N.I. Lazarov modification [1].

We used a borate buffer (6.3g of boric acid and 4g of caustic soda to 1000 ml
of water) with a gradient of potential of 4.7v and the strength of the current
0.2 - 0.3 ma for 1 cm of paper cross section. The electrophoresis took 18 hrs.
The proteinograms were tinted with a dye of the following composition: 0.15g
of bromophenol blue in 100 ml of a 7% solution of trichloroacetic acid.

We obtained clear cut electrophoregrams in which the gamma-globulin fraction
was sometimes divided into 2 subfractions (gamma 1 and gamma 2-globulins). Deter-
mining the quantitative content of the protein fractions was done with the help
of elution and a subsequent colorimetry. The proteinogram was cut into pieces
according to the number of protein fractions. A check of the reproducibility of
the results during a simultaneous investigation of one and the same serum on 7
paper strips demonstrated the sufficient preciseness of the method. The percen-
tage of total protein in the serum was determined by a KLU refractometer.

From studying the sera of 57 donors the following facts were obtained:
Total protein - $7.41 \text{ g\%} \pm 0.07$, albumins - $4.66 \text{ g\%} \pm 0.03$ ($62.9\% \pm 0.62$), globu-
lins: alpha 1 - $0.27 \text{ g\%} \pm 0.01$ ($3.3\% \pm 0.14$), alpha 2 - $0.44 \text{ g\%} \pm 0.01$ ($5.0\% \pm$
 0.18), beta - $0.68 \text{ g\%} \pm 0.02$ ($9.2\% \pm 0.28$), gamma - $1.36 \text{ g\%} \pm 0.04$ ($18.3\% \pm 0.35$),
[gamma 1 - 0.31 g\% (4.2%), gamma 2 - 1.03 g\% (13.9%)] .

In 14 patients with primary and 16 patients with recurrent spotted fever, we
investigated the blood serum 3 times: In the 5-9th day of the febrile period,

○ during the first days of normal temperature, and before discharge. In the event of complications, investigations were conducted more often.

The course of the disease in the cases studied wasn't any different from the clinical aspects of spotted fever, described lately by many researchers. In 6 patients the disease proceeded in a serious form, in 17 it was moderate, and in 7 it was mild. The agglutination reaction with Rickettsia prowazekii was positive in 28 persons (titer 1:40 and higher), and there was a Weil-Felix reaction in 20. The majority of the patients were treated with synthomycin or levomycetin. The average percentage of total protein in the serum during the course of the entire period of observation of the patients was within the limits of normal. During the febrile periods of the disease, dysproteinemia developed which is characterized by hypoalbuminemia, an increase in the percentage of alpha 1 and alpha 2-globulins, a decrease in the level of beta-globulins, and hypergamma-globulinemia. The stated changes were preserved during the first days of normal temperature and the concentration of gamma-globulins increased. Before discharge a tendency was noted toward normalization of the protein content of the blood, but the proteinogram wasn't completely restored. The changes of the gamma 1 and gamma 2-globulins were the same as for the overall gamma-globulins.

Changes of the protein fractions during initial and recurrent spotted fever were analogous during all the periods of the disease. The expressedness of the changes of all the protein fractions, except beta-globulins, was in direct dependence on the gravity of the course of the disease.

During complications (focal pneumonia in 3, pulmonary infarction in 1, and thrombophlebitis of the skin in 1) dysproteinemia developed with the very same trend of changes as during the height of the illness.

○ Hypoalbuminemia and an increase in the level of alpha 2-globulins were especially expressed. In one case we succeeded in observing the stated changes in the

proteinogram one day before the clinical manifestations of focal pneumonia.

The value of the titer of the agglutination reaction with Rickettsia pro-
vazeki didn't depend on the concentration of gamma-globulins, and on the con-
trary. Apparently hypergammaglobulinemia in spotted fever is connected not only
with the accumulation of antibodies but with the increase in the formation of
nonspecific gamma-globulins.

From what has been said it is apparent that the changes in the protein con-
tent of blood serum during spotted fever, just as in other infectious diseases,
are non-specific and stereotyped. Therefore dynamic studies of protein fractions
during spotted fever by paper electrophoresis have only a prognostic significance.
They may be of help in the evaluation of the gravity of the course of the infec-
tious process and in an early diagnosis of complications.

Cited Literature

- (1.) Lazarov, N.I., Labor. Delo, 1957, No. 4, p 11.

Submitted 11/III 1960